

Package: WR (via r-universe)

November 4, 2024

Type Package

Title Win Ratio Analysis of Composite Time-to-Event Outcomes

Version 1.0

Author Lu Mao and Tuo Wang

Maintainer Lu Mao <lmao@biostat.wisc.edu>

URL <https://sites.google.com/view/lmaowisc/>

Description Implements various win ratio methodologies for composite endpoints of death and non-fatal events, including the (stratified) proportional win-fractions (PW) regression models (Mao and Wang, 2020 <[doi:10.1111/biom.13382](https://doi.org/10.1111/biom.13382)>), (stratified) two-sample tests with possibly recurrent nonfatal event, and sample size calculation for standard win ratio test (Mao et al., 2021 <[doi:10.1111/biom.13501](https://doi.org/10.1111/biom.13501)>).

License GPL (>= 2)

Encoding UTF-8

LazyData true

Depends R (>= 3.5.0)

RoxygenNote 7.1.2

Imports survival, cubature, gumbel

Suggests knitr, rmarkdown

VignetteBuilder knitr

NeedsCompilation no

Date/Publication 2021-11-26 21:20:08 UTC

Repository <https://lmaowisc.r-universe.dev>

RemoteUrl <https://github.com/cran/WR>

RemoteRef HEAD

RemoteSha e0837b887272a6a4680beb7e56482c7b6754a3bd

Contents

base	2
gbc	3
gumbel.est	4
hfaction_cpx9	5
non_ischemic	5
plot.pwreg.score	6
print.pwreg	8
print.pwreg.score	8
print.WRrec	9
pwreg	10
score.proc	11
WRrec	12
WRSS	14

Index	16
--------------	-----------

base	<i>Compute the baseline parameters needed for sample size calculation for standard win ratio test</i>
------	---

Description

Compute the baseline parameters ζ_0^2 and δ_0 needed for sample size calculation for standard win ratio test (see [WRSS](#)). The calculation is based on a Gumbel–Hougaard copula model for survival time $D^{(a)}$ and nonfatal event time $T^{(a)}$ for group a (1: treatment; 0: control):

$$P(D^{(a)} > s, T^{(a)} > t) = \exp\left(-\left[\{\exp(a\xi_1)\lambda_D s\}^\kappa + \{\exp(a\xi_2)\lambda_H t\}^\kappa\right]^{1/\kappa}\right),$$

where ξ_1 and ξ_2 are the component-wise log-hazard ratios to be used as effect size in [WRSS](#). We also assume that patients are recruited uniformly over the period $[0, \tau_b]$ and followed until time τ ($\tau \geq \tau_b$), with an exponential loss-to-follow-up hazard λ_L .

Usage

```
base(lambda_D, lambda_H, kappa, tau_b, tau, lambda_L, N = 1000, seed = 12345)
```

Arguments

lambda_D	Baseline hazard λ_D for death.
lambda_H	Baseline hazard λ_H for nonfatal event.
kappa	Gumbel–Hougaard copula correlation parameter κ .
tau_b	Length of the initial (uniform) accrual period τ_b .
tau	Total length of follow-up τ .
lambda_L	Exponential hazard rate λ_L for random loss to follow-up.
N	Simulated sample size for monte-carlo integration.
seed	Seed for monte-carlo simulation.

Value

A list containing real number ζ_0^2 for ζ_0^2 and bivariate vector δ_0 for δ_0 .

References

Mao, L., Kim, K. and Miao, X. (2021). Sample size formula for general win ratio analysis. *Biometrics*, <https://doi.org/10.1111/biom.13501>.

See Also

[gumbel.est](#), WRSS

Examples

```
# see the example for WRSS
```

gbc

A subset of the German Breast Cancer study data

Description

These are a subset of the German Breast Cancer study data.

Usage

```
gbc
```

Format

A data frame with 985 rows and 12 variables:

id subject IDs

time event times (months)

status event status; 0:censoring, 1:death, 2:cancer recurrence

hormone treatment indicator: 1=Hormone therapy; 2=standard therapy

age age at diagnosis (years)

menopause menopausal Status; 1=No; 2=Yes

size tumor size

grade tumor grade, 1-3

nodes number of nodes involved

prog_recp number of progesterone receptors

estrg_recp number of estrogen receptors

References

Sauerbrei, W., Royston, P., Bojar, H., Schmoor, C. and Schumacher, M. (1999). Modelling the effects of standard prognostic factors in node-positive breast cancer. German Breast Cancer Study Group (GBSG). *British Journal of Cancer*, 79, 1752–1760.

Hosmer, D.W. and Lemeshow, S. and May, S. (2008) *Applied Survival Analysis: Regression Modeling of Time to Event Data: Second Edition*, John Wiley and Sons Inc., New York, NY

gumbel.est	<i>Estimate baseline parameters in the Gumbel–Hougaard model for sample size calculation using pilot data</i>
------------	---

Description

Estimate baseline parameters in the Gumbel–Hougaard model described in [base](#) for sample size calculation using pilot study data.

Usage

```
gumbel.est(id, time, status)
```

Arguments

id	A vector of unique patient identifiers.
time	A numeric vector of event times.
status	A vector of event type variable; 2 = nonfatal event, 1 = death, and 0 = censoring.

Value

A list containing `lambda_D` for λ_D , `lambda_H` for λ_H , and `kappa` for κ in the Gumbel–Hougaard model.

References

Mao, L., Kim, K. and Miao, X. (2021). Sample size formula for general win ratio analysis. *Biometrics*, <https://doi.org/10.1111/biom.13501>.

See Also

[base](#), [WRSS](#)

Examples

```
# see the example for WRSS
```

hfaction_cpx9	<i>A subset of the HF-ACTION study data on high-risk non-ischemic heart failure patients</i>
---------------	--

Description

These are data on a subgroup of 426 high-risk non-ischemic patients in the HF-ACTION study.

Usage

hfaction_cpx9

Format

A data frame with 1,448 rows and 5 variables:

patid patient ID

time event times (months)

status event status; 0:censoring, 1:death, 2:hospitalization

trt_ab treatment indicator: 1: exercise training, 0: usual care

age60 1: 60 years or older, 0: less than 60 years old

References

O'Connor, C. M., Whellan, D. J., Lee, K. L., Keteyian, S. J., Cooper, L. S., Ellis, S. J., Leifer, E. S., Kraus, W. E., Kitzman, D. W., Blumenthal, J. A. et al. (2009). Efficacy and safety of exercise training in patients with chronic heart failure: HF-ACTION randomized controlled trial. *Journal of the American Medical Association*, 301, 1439–1450.

non_ischemic	<i>A subset of the HF-ACTION study data on non-ischemic heart failure patients with full covariate measurement.</i>
--------------	---

Description

These are a subset of the data on 451 non-ischemic patients in the HF-ACTION study will complete baseline covariates.

Usage

non_ischemic

Format

A data frame with 751 rows and 16 variables:

ID subject IDs

time event times (days)

status event status; 0:censoring, 1:death, 2:hospitalization

trt_ab treatment indicator: 1=exercise training; 0=usual care

age patient age in years

sex 1=female; 2=male

Black.vs.White 1=black; 0=otherwise

Other.vs.White 1=race other than black or white; 0=otherwise

bmi body mass index

bipllvf (biplane) left-ventricular ejection fraction

hyperten indicator for history of hypertension

COPD indicator for history of COPD

diabetes indicator for history of diabetes

acei indicator for current use of ACE inhibitors

betab indicator for current use of beta blockers

smokecurr indicator for current smoker

References

O'Connor, C. M., Whellan, D. J., Lee, K. L., Keteyian, S. J., Cooper, L. S., Ellis, S. J., Leifer, E. S., Kraus, W. E., Kitzman, D. W., Blumenthal, J. A. et al. (2009). Efficacy and safety of exercise training in patients with chronic heart failure: HF-ACTION randomized controlled trial. *Journal of the American Medical Association*, 301, 1439–1450.

plot.pwreg.score

Plot the standardized score processes

Description

Plot the standardized score processes.

Usage

```
## S3 method for class 'pwreg.score'
plot(
  x,
  k,
  xlab = "Time",
  ylab = "Standardized score",
```

```
    lty = 1,  
    frame.plot = TRUE,  
    add = FALSE,  
    ylim = c(-3, 3),  
    xlim = NULL,  
    lwd = 1,  
    ...  
)
```

Arguments

<code>x</code>	an object of class <code>pwreg.score</code> .
<code>k</code>	A positive integer indicating the order of covariate to be plotted. For example, <code>k=3</code> requests the standardized score process for the third covariate in the covariate matrix <code>Z</code> .
<code>xlab</code>	a title for the x axis.
<code>ylab</code>	a title for the y axis.
<code>lty</code>	the line type. Default is 1.
<code>frame.plot</code>	a logical variable indicating if a frame should be drawn in the 1D case.
<code>add</code>	a logical variable indicating whether add to current plot?
<code>ylim</code>	a vector indicating the range of y-axis. Default is (-3,3).
<code>xlim</code>	a vector indicating the range of x-axis. Default is NULL.
<code>lwd</code>	the line width, a positive number. Default is 1.
<code>...</code>	further arguments passed to or from other methods

Value

A plot of the standardized score process for object `pwreg.score`.

See Also

[score.proc](#)

Examples

```
# see the example for score.proc
```

print.pwreg *Print the results of the proportional win-fractions regression model*

Description

Print the results of the proportional win-fractions regression model.

Usage

```
## S3 method for class 'pwreg'  
print(x, ...)
```

Arguments

x an object of class pwreg.
... further arguments passed to or from other methods

Value

Print the results of pwreg object

See Also

[pwreg](#)

Examples

```
# see the example for pwreg
```

print.pwreg.score *Print information on the content of the pwreg.score object*

Description

Print information on the content of the pwreg.score object

Usage

```
## S3 method for class 'pwreg.score'  
print(x, ...)
```

Arguments

x A object of class pwreg.score.
... further arguments passed to or from other methods.

Value

Print the results of `pwreg.score` object.

See Also

[score.proc](#)

Examples

```
# see the example for score.proc
```

<code>print.WRrec</code>	<i>Print the results of the two-sample recurrent-event win ratio analysis</i>
--------------------------	---

Description

Print the results of the two-sample recurrent-event win ratio analysis.

Usage

```
## S3 method for class 'WRrec'  
print(x, ...)
```

Arguments

<code>x</code>	an object of class <code>WRrec</code> .
<code>...</code>	further arguments passed to or from other methods.

Value

Print the results of `WRrec` object.

See Also

[WRrec](#)

Examples

```
# see the example for WRrec
```

pwreg

Fit a standard proportional win-fractions (PW) regression model

Description

Fit a standard proportional win-fractions (PW) regression model.

Usage

```
pwreg(
  ID,
  time,
  status,
  Z,
  rho = 0,
  strata = NULL,
  fixedL = TRUE,
  eps = 1e-04,
  maxiter = 50
)
```

Arguments

ID	a vector of unique subject-level identifiers.
time	a vector of event times.
status	a vector of event type labels. 0: censoring, 1:death and 2: non-fatal event.
Z	a matrix or a vector of covariates.
rho	a non-negative number as the power of the survival function used in the weight. Default (rho=0) is recommended. If there is a 'strata' argument, then 'rho' is ignored.
strata	a vector of stratifying variable if a stratified model is desired.
fixedL	logical variable indicating which variance estimator to be used. If 'TRUE', the type I variance estimator (for a small number strata) is used; otherwise the type II variance estimator (for a large number strata) is used.
eps	precision for the convergence of Newton-Raphson algorithm.
maxiter	maximum number of iterations allow for the Newton-Raphson algorithm.

Value

An object of class pwreg with the following components. beta:a vector of estimated regression coefficients. Var:estimated covariance matrix for beta. conv: boolean variable indicating whether the algorithm converged within the maximum number of iterations.

References

- Mao, L. and Wang, T. (2020). A class of proportional win-fractions regression models for composite outcomes. *Biometrics*, 10.1111/biom.13382
- Wang, T. and Mao, L. (2021+). Stratified Proportional Win-fractions Regression Analysis.

See Also

[score.proc](#), [print.pwreg](#)

Examples

```
library(WR)
head(non_ischemic)
id_unique <- unique(non_ischemic$ID)

# Randomly sample 200 subjects from non_ischemic data
set.seed(2019)
id_sample <- sample(id_unique, 200)
non_ischemic_reduce <- non_ischemic[non_ischemic$ID %in% id_sample, ]

# Use the reduced non_ischemic data for analysis
nr <- nrow(non_ischemic_reduce)
p <- ncol(non_ischemic_reduce)-3
ID <- non_ischemic_reduce[, "ID"]
time <- non_ischemic_reduce[, "time"]
status <- non_ischemic_reduce[, "status"]
Z <- as.matrix(non_ischemic_reduce[, 4:(3+p)], nr, p)
## unstratified analysis
pwreg.obj <- pwreg(time=time, status=status, Z=Z, ID=ID)
print(pwreg.obj)
## Not run:
## stratified PW by sex
sex <- Z[, 3]
## take out sex from the covariate matrix
Z1 <- Z[, -3]
pwreg.obj1 <- pwreg(time=time, status=status, Z=Z1, ID=ID, strata=sex)
print(pwreg.obj1)

## End(Not run)
```

score.proc

Computes the standardized score processes

Description

Computes the standardized score processes for the covariates.

Usage

```
score.proc(obj, t = NULL)
```

Arguments

<code>obj</code>	an object of class <code>pwreg</code> .
<code>t</code>	a vector containing times. If not specified, the function will use all unique event times from the data.

Value

An object of class `pwreg.score` consisting of `t`: a vector of times; and `score`: a matrix whose rows are the standardized score processes as a function of `t`.

References

Mao, L. and Wang, T. (2020). A class of proportional win-fractions regression models for composite outcomes. *Biometrics*, 10.1111/biom.13382

See Also

[pwreg](#), [print.pwreg](#)

Examples

```
library(WR)
head(non_ischemic)

# Randomly sample 200 subjects from non_ischemic data
id_unique <- unique(non_ischemic$ID)
set.seed(2019)
id_sample <- sample(id_unique, 200)
non_ischemic_reduce <- non_ischemic[non_ischemic$ID %in% id_sample, ]

# Use the reduced non_ischemic data for analysis
nr <- nrow(non_ischemic_reduce)
p <- ncol(non_ischemic_reduce)-3
ID <- non_ischemic_reduce[, "ID"]
time <- non_ischemic_reduce[, "time"]
status <- non_ischemic_reduce[, "status"]
Z <- as.matrix(non_ischemic_reduce[, 4:(3+p)], nr, p)
pwreg.obj <- pwreg(time=time, status=status, Z=Z, ID=ID)
score.obj <- score.proc(pwreg.obj)
#plot the standardized score process for the first covariate
plot(score.obj, k = 1)
```

Description

Perform stratified two-sample test of possibly recurrent nonfatal event and death using the recommended last-event assisted win ratio (LWR), and/or naive win ratio (NWR) and first-event assisted win ratio (FWR) (Mao et al., 2022). The LWR and FWR reduce to the standard win ratio of Pocock et al. (2012).

Usage

```
WRrec(ID, time, status, trt, strata = NULL, naive = FALSE)
```

Arguments

ID	A vector of unique patient identifiers.
time	A numeric vector of event times.
status	A vector of event type variable; 2 = recurrent event, 1 = death, and 0 = censoring.
trt	A vector of binary treatment indicators.
strata	A vector of categorical variable for strata; Default is NULL, which leads to unstratified analysis.
naive	If TRUE, results for NWR and FWR will be provided in addition to LWR; Default is FALSE, which gives LWR only.

Value

An object of class WRrec, which contains the following elements.

theta	A bivariate vector of win/loss fractions by LWR.
log.WR, se	Log-win ratio estimate and its standard error by LWR.
pval	p -value by the LWR test.
theta.naive	A bivariate vector of win/loss fractions by NWR.
log.WR.naive, se.naive	Log-win ratio estimate and its standard error by NWR.
theta.FI	A bivariate vector of win/loss fractions by FWR.
log.WR.FI, se.FI	Log-win ratio estimate and its standard error by FWR.
...	

References

- Mao, L., Kim, K. and Li, Y. (2022). On recurrent-event win ratio. *Statistical Methods in Medical Research*, under review.
- Pocock, S., Ariti, C., Collier, T., and Wang, D. (2012). The win ratio: a new approach to the analysis of composite endpoints in clinical trials based on clinical priorities. *European Heart Journal*, 33, 176–182.

See Also

[print.WRrec.](#)

Examples

```
## load the HF-ACTION trial data
library(WR)
head(hfaction_cpx9)
dat<-hfaction_cpx9
## Comparing exercise training to usual care by LWR, FWR, and NWR
obj<-WRrec(ID=dat$patid,time=dat$time,status=dat$status,
           trt=dat$trt_ab,strata=dat$age60,naive=TRUE)
## print the results
obj
```

WRSS

Compute the sample size for standard win ratio test

Description

Compute the sample size for standard win ratio test.

Usage

```
WRSS(xi, bparam, q = 0.5, alpha = 0.05, side = 2, power = 0.8)
```

Arguments

xi	A bivariate vector of hypothesized component-wise (treatment-to-control) log-hazard ratios under the Gumbel–Hougaard copula model described in base .
bparam	A list containing baseline parameters zeta2 for ζ_0^2 and delta for δ_0 ; Can directly use the output of base .
q	Proportion of patients assigned to treatment.
alpha	Type I error rate.
side	2-sided or 1-sided test.
power	Target power.

Value

A list containing n, the computed sample size.

References

Mao, L., Kim, K. and Miao, X. (2021). Sample size formula for general win ratio analysis. *Biometrics*, <https://doi.org/10.1111/biom.13501>.

See Also

[gumbel.est, base](#)

Examples

```
# The following is not run in package checking to save time.
## Not run:
## load the package and pilot dataset
library(WR)
head(hfaction_cpx9)
dat<-hfaction_cpx9
## subset to control group
pilot<-dat[dat$trt_ab==0,]

## get the data ready for gumbel.est()
id<-pilot$patid
## convert time from month to year
time<-pilot$time/12
status<-pilot$status
## compute the baseline parameters for the Gumbel--Hougaard
## copula for death and hospitalization
gum<-gumbel.est(id, time, status)

## get the baseline parameters
lambda_D<-gum$lambda_D
lambda_H<-gum$lambda_H
kappa<-gum$kappa
## set up design parameters and use base()
## to calculate bparam for WRSS()
# max follow-up 4 years
tau<-4
# 3 years of initial accrual
tau_b<-3
# loss to follow-up rate
lambda_L=0.05
# compute the baseline parameters
bparam<-base(lambda_D,lambda_H,kappa,tau_b,tau,lambda_L)
bparam

## sample size with power=0.8 under hazard ratios
## 0.9 and 0.8 for death and hospitalization, respectively.
WRSS(xi=log(c(0.9,0.8)),bparam=bparam,q=0.5,alpha=0.05,
      power=0.8)$n
## sample size under the same set-up but with power 0.9
WRSS(xi=log(c(0.9,0.8)),bparam=bparam,q=0.5,alpha=0.05,
      power=0.9)$n

## End(Not run)
```

Index

- * **WRSS**
 - base, [2](#)
 - gumbel.est, [4](#)
 - WRSS, [14](#)
- * **WRrec**
 - print.WRrec, [9](#)
 - WRrec, [12](#)
- * **datasets**
 - gbc, [3](#)
 - hfaction_cpx9, [5](#)
 - non_ischemic, [5](#)
- * **pwreg**
 - plot.pwreg.score, [6](#)
 - print.pwreg, [8](#)
 - print.pwreg.score, [8](#)
 - pwreg, [10](#)
 - score.proc, [11](#)

base, [2](#), [4](#), [14](#), [15](#)

gbc, [3](#)

gumbel.est, [3](#), [4](#), [15](#)

hfaction_cpx9, [5](#)

non_ischemic, [5](#)

plot.pwreg.score, [6](#)

print.pwreg, [8](#), [11](#), [12](#)

print.pwreg.score, [8](#)

print.WRrec, [9](#), [14](#)

pwreg, [8](#), [10](#), [12](#)

score.proc, [7](#), [9](#), [11](#), [11](#)

WRrec, [9](#), [12](#)

WRSS, [2-4](#), [14](#)